

STN Search History

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTAGXP1614

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	MAY 01	New CAS web site launched
NEWS	3	MAY 08	CA/CAPLUS Indian patent publication number format defined
NEWS	4	MAY 14	RDISCLOSURE on STN Easy enhanced with new search and display fields
NEWS	5	MAY 21	BIOSIS reloaded and enhanced with archival data
NEWS	6	MAY 21	TOXCENTER enhanced with BIOSIS reload
NEWS	7	MAY 21	CA/CAPLUS enhanced with additional kind codes for German patents
NEWS	8	MAY 22	CA/CAPLUS enhanced with IPC reclassification in Japanese patents
NEWS	9	JUN 27	CA/CAPLUS enhanced with pre-1967 CAS Registry Numbers
NEWS	10	JUN 29	STN Viewer now available
NEWS	11	JUN 29	STN Express, Version 8.2, now available
NEWS	12	JUL 02	LEMBASE coverage updated
NEWS	13	JUL 02	LMEDLINE coverage updated
NEWS	14	JUL 02	SCISEARCH enhanced with complete author names
NEWS	15	JUL 02	CHEMCATS accession numbers revised
NEWS	16	JUL 02	CA/CAPLUS enhanced with utility model patents from China
NEWS	17	JUL 16	CAPLUS enhanced with French and German abstracts
NEWS	18	JUL 18	CA/CAPLUS patent coverage enhanced
NEWS	19	JUL 26	USPATFULL/USPAT2 enhanced with IPC reclassification
NEWS	20	JUL 30	USGENE now available on STN
NEWS	21	AUG 06	CAS REGISTRY enhanced with new experimental property tags
NEWS	22	AUG 06	BEILSTEIN updated with new compounds
NEWS	23	AUG 06	FSTA enhanced with new thesaurus edition
NEWS	24	AUG 13	CA/CAPLUS enhanced with additional kind codes for granted patents
NEWS	25	AUG 20	CA/CAPLUS enhanced with CAS indexing in pre-1907 records
NEWS	26	AUG 27	Full-text patent databases enhanced with predefined patent family display formats from INPADOCDB
NEWS	27	AUG 27	USPATOLD now available on STN
NEWS	28	AUG 28	CAS REGISTRY enhanced with additional experimental spectral property data
NEWS EXPRESS	29	JUNE 2007:	CURRENT WINDOWS VERSION IS V8.2, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 05 JULY 2007.
NEWS HOURS			STN Operating Hours Plus Help Desk Availability
NEWS LOGIN			Welcome Banner and News Items
NEWS IPC8			For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 15:51:23 ON 28 AUG 2007

=> fil reg

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 15:51:30 ON 28 AUG 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 27 AUG 2007 HIGHEST RN 945649-99-0

DICTIONARY FILE UPDATES: 27 AUG 2007 HIGHEST RN 945649-99-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007


Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10597663.str



chain nodes :
 26 28 31 34 38 39 42 44
 ring nodes :
 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23
 24 25 27 29 30 32 33 35 36 37 40 41 43
 chain bonds :
 25-26 27-28 30-31 32-34 36-38 37-39 41-42 43-44
 ring bonds :
 1-2 1-6 2-3 3-4 3-25 4-5 5-6 5-30 7-8 7-12 7-29 8-9 9-10 10-11 10-40
 11-12 13-14 13-18 13-33 14-15 15-16 16-17 16-35 17-18 19-20 19-24 20-21
 20-43 21-22 22-23 23-24 24-37 25-27 27-29 30-32 32-33 35-36 36-37 40-41
 41-43
 exact/norm bonds :
 3-25 5-30 7-29 10-40 13-33 16-35 20-43 24-37 25-26 25-27 27-29 30-31
 30-32 32-33 35-36 36-37 37-39 40-41 41-43 43-44
 exact bonds :
 27-28 32-34 36-38 41-42
 normalized bonds :
 1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12 13-14 13-18 14-15
 15-16 16-17 17-18 19-20 19-24 20-21 21-22 22-23 23-24


Match level :
 1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
 11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom
 20:Atom 21:Atom 22:Atom 23:Atom 24:Atom 25:Atom 26:CLASS 27:Atom 28:CLASS
 29:Atom 30:Atom 31:CLASS 32:Atom 33:Atom 34:CLASS 35:Atom 36:Atom 37:Atom
 38:CLASS 39:CLASS 40:Atom 41:Atom 42:CLASS 43:Atom 44:CLASS

L1 STRUCTURE UPLOADED

=> d 11

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 15:52:04 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 1231 TO ITERATE

100.0% PROCESSED 1231 ITERATIONS

9 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 22516 TO 26724

PROJECTED ANSWERS: 9 TO 360

L2 9 SEA SSS SAM L1



=> d scan

L2 9 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN


IN Glycine, N-(diphenylacetyl)glycyl-, [[(4-methylphenyl)sulfonyl]sulfinimido
yl]di-3,1-propanediyl ester, rotaxane compd. with 3,11,18,26-
tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-
5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone
(1:1) (9CI)

MF C49 H53 N5 O10 S2 . C32 H28 N4 O4

CM 1



CM 2




HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 9 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
 IN Copper, [μ -[N,N'-bis[2-[bis[(2-pyridinyl- κ N)methyl]amino- κ N]ethyl]-2-butenediamide]]tetrachlorodi-, rotaxane compd. with
 N,N-dimethylformamide and 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9
 .120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-
 4,10,19,25-tetrone (1:2:1) (9CI)
 MF C32 H36 Cl4 Cu2 N8 O2 . C32 H28 N4 O4 . 2 C3 H7 N O

CM 1


PAGE 1-A




PAGE 2-A



CM 2




CM 3



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L2 9 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
IN Carbamic acid, [22-(1,1-dimethylethyl)-4,10,19,25-tetraoxo-3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(34),13,15,20,22,24(33),28,30,31,35-dodecaen-7-yl]-, 2-hexadecenyl ester (9CI)
MF C53,H67 N5 O6
CI COM



—Bu-t


PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 9 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
IN L-Proline, N-(diphenylacetyl)glycyl-, 2,2-diphenylethyl ester, rotaxane compd. with 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:1) (9CI)
MF C35 H34 N2 O4 . C32 H28 N4 O4

CM 1

Absolute stereochemistry.




CM 2




HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L2 9 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
 IN Glycine, N-[4-[(10-mercaptodecyl)oxy]-3,5-dimethoxybenzoyl]glycyl-,
 2,2-diphenylethyl ester, rotaxane compd. with 3,11,18,26-
 tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-
 5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone
 (1:1) (9CI)
 MF C37 H48 N2 O7 S . C32 H28 N4 O4

CM 1



CM 2



HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 9 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN

IN Butanediamide, N-[11-[(2,2-diphenylethyl)amino]-11-oxoundecyl]-N'-[2-(2'H-[5,6]fullereno-C60-Ih-[1,9-c]pyrrol-1'(5'H)-yl)ethyl]-, radical ion(1-), rotaxane compd. with 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:1)


MF C93 H48 N4 O3 . C32 H28 N4 O4


CM 1

PAGE 1-A




PAGE 1-B






CM 2



HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 9 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
 IN Glycinamide, N,N-bis(2-pyridinylmethyl)glycyl-N-[12-[[4-(2,2-diphenylethoxy)-1,4-dioxobutyl]amino]dodecyl]-, rotaxane compd. with 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:1) (9CI)
 MF C46 H60 N6 O5 . C32 H28 N4 O4

CM 1



—CH₂—CHPh₂


CM 2



HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L2 9 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
 IN 2-Butenediamide, N,N,N',N'-tetrakis[6-[[(hexahydro-2-oxo-1H-azepin-1-yl)carbonyl]amino]hexyl]-, (2E)-, rotaxane compd. with
 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-
 5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone
 (1:1), polymer with α -(2-aminomethylethyl)- ω -(2-aminomethylethoxy)poly[oxy(methyl-1,2-ethanediyl)] (9CI)
 MF (C₅₆ H₉₄ N₁₀ O₁₀ . C₃₂ H₂₈ N₄ O₄ . (C₃ H₆ O)_n C₆ H₁₆ N₂ O)_x
 CI PMS

CM 1





2 (D1—Me)


CM 2

CM 3

Double bond geometry as shown.



CM 4




HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L2 9 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
 IN L-Methionine, N-(diphenylacetyl)glycyl-, 10-[3,5-bis(1,1-dimethylethyl)phenoxy]decyl ester, rotaxane compd. with
 7,22-bis(2-propenyloxy)-3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.1
 20,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-
 4,10,19,25-tetrone (1:1) (9CI)


MF C45 H64 N2 O5 S . C38 H36 N4 O6

CM 1

Absolute stereochemistry.



CM 2



ALL ANSWERS HAVE BEEN SCANNED

=> 1

FILE 'CAPLUS' ENTERED AT 15:57:13 ON 28 AUG 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 28 Aug 2007 VOL 147 ISS 10

FILE LAST UPDATED: 27 Aug 2007 (20070827/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=>

=> d ibib abs hitstr 1-

YOU HAVE REQUESTED DATA FROM 8 ANSWERS - CONTINUE? Y/(N):y

L6 ANSWER 1 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:1042566 CAPLUS Full-text

DOCUMENT NUMBER: 146:7581

TITLE: Reverse Shuttling in a Fullerene-Stoppered Rotaxane

AUTHOR(S): Mateo-Alonso, Aurelio; Fioravanti, Giulia; Marcaccio, Massimo; Paolucci, Francesco; Jagesar, Dhiredj C.; Brouwer, Albert M.; Prato, Maurizio

CORPORATE SOURCE: INSTM unit of Trieste and Dipartimento di Scienze Farmaceutiche, Universita degli Studi di Trieste, Trieste, 34127, Italy

SOURCE: Organic Letters (2006), 8(22), 5173-5176

CODEN: ORLEF7; ISSN: 1523-7060

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 146:7581

AB The prepn. and characterization of a solvent-switchable rotaxane that shuttles in the opposite direction to that expected are reported. The reverse shuttling is confirmed by NMR spectroscopy and can be monitored by cyclic voltammetry. The electrochem. generated anions on the fullerene moiety are stabilized by the closer proximity of the macrocycle.

IT 913943-50-7

RL: FMU (Formation, unclassified); PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)
(reverse shuttling in fullerene-stoppered rotaxane)

RN 913943-50-7 CAPLUS

CN Butanediamide, N-[11-[(2,2-diphenylethyl)amino]-11-oxoundecyl]-N'-[2-(2'H-[5,6]fullereno-C60-Ih-[1,9-c]pyrrol-1'(5'H)-yl)ethyl]-, radical ion(1-), rotaxane compd. with 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:1) (CA INDEX NAME)


CM 1



CRN 913943-49-4

CMF C93 H48 N4 O3

CCI RIS


PAGE 1-A





CM 2


CRN 169203-75-2
CMF C32 H28 N4 O4




REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 2 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2006:219952 CAPLUS Full-text
DOCUMENT NUMBER: 144:444544
TITLE: An allosterically regulated molecular shuttle
AUTHOR(S): Marlin, Dana S.; Cabrera, Diego Gonzalez; Leigh, David A.; Slawin, Alexandra M. Z.
CORPORATE SOURCE: School of Chemistry, University of Edinburgh, Edinburgh, EH9 3JJ, UK
SOURCE: Angewandte Chemie, International Edition (2006), 45(9), 1385-1390
CODEN: ACIEF5; ISSN: 1433-7851
PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 144:444544
AB A switchable rotaxane was developed in which the binding of a metal ion (e.g. Cd²⁺) to a bis(picoly)amine fragment that acts as one of the stoppers induces a shift of the macrocyclic component along the thread from one binding site to another weaker binding site. Crystal structures were determined for two metal-free rotaxanes and for three copper complexes.
IT 884592-09-0P
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (preparation and crystal structure of copper complex of a rotaxane with thread having dipicolylamine chelating group)
RN 884592-09-0 CAPLUS
CN Copper, [μ -[N,N'-bis[2-[bis[(2-pyridinyl- κ N)methyl]amino- κ N]ethyl]-2-butenediamide]]tetrachlorodi-, rotaxane compd. with N,N-dimethylformamide and 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:2:1) (9CI) (CA INDEX NAME)
CM 1
CRN 884592-08-9
CMF C32 H36 Cl4 Cu2 N8 O2
CCI CCS

PAGE 1-A






CM 2

CRN 169203-75-2


CMF C32 H28 N4 O4



CM 3

CRN 68-12-2

CMF C3 H7 N O



REFERENCE COUNT: 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 3 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2006:17634 CAPLUS Full-text
 DOCUMENT NUMBER: 144:245942
 TITLE: Complexation-induced translational isomerism:
 Shuttling through stepwise competitive binding
 AUTHOR(S): Marlin, Dana S.; Cabrera, Diego Gonzalez; Leigh, David
 A.; Slawin, Alexandra M. Z.
 CORPORATE SOURCE: School of Chemistry, University of Edinburgh,
 Edinburgh, EH93JJ, UK
 SOURCE: Angewandte Chemie, International Edition (2006),
 45(1), 77-83
 CODEN: ACIEF5; ISSN: 1433-7851
 PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA

DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 144:245942

AB Progressive binding of a transition-metal ion to a peptide station displaces the macrocycle to an alternative station in a H-bonded mol. shuttle. Cu(HL)Cl₂ (HL = 5-((RCH₂)₂NCH₂CONH)C₆H₃(CO₂Me)₂-1,3 (R = 2-pyridyl)), was prepared and reacted with CuCl₂ or NaH to give [Cu(HL)Cl]₂[CuCl₄] and CuLCl, resp. (RCH₂)₂NCH₂CONHCH₂CONHCH₂CHPh₂ (H₂L₁) reacted with Cu and Cd salts to give Cu(H₂L₁)Cl₂ and Cd(H₂L₁)(NO₃)₂ which were deprotonated to give Cu(HL₁)Cl and Cd(HL₁)(NO₃), resp., with change in the coordination environment. The rotaxane H₂L₁.L₂ (L = 2,6,15,19-tetraoxo-1,7,14,20-tetraaza-3,5,9,12,16,18,22,25-tetrabenzocyclohexancosane), prepared from p-xylylenediamine and isophthaloyl chloride, was prepared and reacted with Cu and Cd salts to give Cu(H₂L₁.L₂)Cl₂ and Cd(H₂L₁.L₂)(NO₃)₂, resp. Deprotonation of Cu(H₂L₁.L₂)Cl₂ gave Cu(HL₁.L₂)Cl with change in the coordination environment and subsequently with degradation of the rotaxane structure. The rotaxane H₂L₃.L₂ (H₂L₃ = (RCH₂)₂NCH₂CONHCH₂CONH(CH₂)₁₂NHCOCH₂CH₂CO₂CH₂CHPh₂) reacted with Cu and Cd salts to give Cu(H₂L₃.L₂)Cl₂ and Cd(H₂L₃.L₂)(NO₃)₂ which were deprotonated to give Cu(HL₃.L₂)Cl and Cd(HL₃.L₂)(NO₃), resp., with shuttling of L₂ to an alternative station in the H-bonded mol. shuttle. The crystal structures of several of these complexes were determined

IT 876289-70-2

RL: RCT (Reactant); RACT (Reactant or reagent)
 (rotaxane; preparation and complexation with cadmium and copper)


RN 876289-70-2 CAPLUS

CN Glycinamide, N,N-bis(2-pyridinylmethyl)glycyl-N-[12-[[4-(2,2-diphenylethoxy)-1,4-dioxobutyl]amino]dodecyl]-, rotaxane compd. with 3,11,18,26-tetraazapentacyclo[26.2.2.2¹³.16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 876289-69-9

CMF C46 H60 N6 O5




PAGE 1-B

— CH₂ — CHPh₂

CM 2

CRN 169203-75-2

CMF C32 H28 N4 O4



REFERENCE COUNT: 98 THERE ARE 98 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 4 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:638951 CAPLUS Full-text

DOCUMENT NUMBER: 139:292587

TITLE: Rotaxane building blocks bearing blocked isocyanate stoppers: Polyrotaxanes through post-assembly chain extension

AUTHOR(S): Kidd, Timothy J.; Loontjens, Ton J. A.; Leigh, David A.; Wong, Jenny K. Y.

CORPORATE SOURCE: DSM Research, Geleen, 6160 MD, Neth.

SOURCE: Angewandte Chemie, International Edition* (2003), 42(29), 3379-3383

CODEN: ACIEF5; ISSN: 1433-7851

PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A simple, versatile route to polyamide-polyoxyalkylene-polyurea polyrotaxanes with a variety of topologies is available through the synthesis and subsequent polymerization of readily accessible, stable, rotaxane monomers containing bulky, blocked isocyanate stopper groups. These monomeric [2]rotaxanes represent a new kind of functional-chain extender that is wholly compatible with methodologies currently used to make a range of com. polymers and could thus lead to the facile incorporation of a number of rotaxane systems into materials where the effects of the mech. bond could be tested and realistically exploited. In principle, this system can also be applied to the modular generation of rotaxane-based polymeric materials or devices in which individual components can be introduced in different combinations through "bottom-up" processing.

IT 609842-48-0P

RL: SPN (Synthetic preparation); PREP (Preparation)

(rotaxane building blocks bearing blocked isocyanate stoppers used to prepare polyrotaxanes through post-assembly chain extension)

RN 609842-48-0 CAPLUS


CN 2-Butenediamide, N,N,N',N'-tetrakis[6-[[[(hexahydro-2-oxo-1H-azepin-1-yl)carbonyl]amino]hexyl]-, (2E)-, rotaxane compd. with 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:1), polymer with α -(2-aminomethylethyl)- ω -(2-aminomethylethoxy)poly[oxy(methyl-1,2-ethanediyl)] (9CI) (CA INDEX NAME)

CM 1

CRN 9046-10-0

CMF (C3 H6 O)_n C6 H16 N2 O

CCI IDS, PMS



2 (D1-Me)

CM 2

CRN 609842-43-5

CMF C56 H94 N10 O10 . C32 H28 N4 O4


CM 3

CRN 609842-42-4


CMF C56 H94 N10 O10

Double bond geometry as shown.

PAGE 1-A




PAGE 1-B



CM 4

CRN 169203-75-2
CMF C32 H28 N4 O4



REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 5 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:159908 CAPLUS Full-text

DOCUMENT NUMBER: 136:340989

TITLE: Switching "On" and "Off" the Expression of Chirality in Peptide Rotaxanes

AUTHOR(S): Asakawa, Masumi; Brancato, Giuseppe; Fanti, Marianna; Leigh, David A.; Shimizu, Toshimi; Slawin, Alexandra M. Z.; Wong, Jenny K. Y.; Zerbetto, Francesco; Zhang, Songwei

CORPORATE SOURCE: Nanoarchitectonics Research Center, National Institute of Advanced Industrial Science and Technology, Tsukuba, Ibaraki, 305-8565, Japan

SOURCE: Journal of the American Chemical Society (2002), 124(12), 2939-2950
CODEN: JACSAT; ISSN: 0002-7863

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 136:340989

AB The hydrogen-bond-directed synthesis, x-ray crystal structures, and optical properties of the first chiral peptide rotaxanes are reported. Collectively these systems provide the first examples of single mol. species where the expression of chirality in the form of a CD response can selectively be switched "on" or "off", and its magnitude altered, through controlling the interactions between mech. interlocked submol. components. The switching is achievable both thermally and through changes in the nature of the environment. Peptido[2]rotaxanes consisting of an intrinsically achiral benzylic amide macrocycle locked onto various chiral dipeptide (Gly-Ala, Gly-Leu, Gly-Met, Gly-Phe, and Gly-Pro) threads exhibit strong (10-20k deg cm² dmol⁻¹) neg. induced CD (θ) values in nonpolar solvents (e.g. CHCl₃), where the intramol. hydrogen bonding between thread and macrocycle is maximized. In polar solvents (e.g., MeOH), where the intercomponent hydrogen bonding is weakened, or switched off completely, the elliptical polarization falls close to zero in some cases and can even be switched to large pos. values in others. Importantly, the mechanism of generating the switchable CD response in the chiral peptide rotaxanes is also determined: a combination of semiempirical calcns. and geometrical modeling using the continuous chirality measure (CCM) shows that the chirality is transmitted from the amino acid asym. center on the thread via the macrocycle to the C-terminal stopper of the rotaxane. This

understanding could have important implications for other areas where chiral transmission from one chemical entity to another underpins a phys. or chemical response, such as the seeding of supertwisted nematic liquid crystalline phases or asym. synthesis.

IT 419537-14-7P

RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)

(preparation, crystal structures and optical properties of peptide rotaxanes

to study the switching "on" and "off" their expression of chirality)

RN 419537-14-7 CAPLUS


CN L-Proline, N-(diphenylacetyl)glycyl-, 2,2-diphenylethyl ester, rotaxane compd. with 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 419537-13-6

CMF C35 H34 N2 O4


Absolute stereochemistry.



CM 2

CRN 169203-75-2

CMF C32 H28 N4 O4



REFERENCE COUNT:

75

THERE ARE 75 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 6 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2001:21352 CAPLUS Full-text


DOCUMENT NUMBER: 134:72032

TITLE: Intermediates for manufacture of polymers containing rotaxanes having a cyclotetraamide ring in main chains

INVENTOR(S): Asakawa, Masumi; Murphy, Aden; Leigh, David A.;

Shimizu, Toshimi
 PATENT ASSIGNEE(S): Agency of Industrial Sciences and Technology, Japan;
 National Institute of Advanced Industrial Science and
 Technology
 SOURCE: Jpn. Kokai Tokkyo Koho, 15 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001002663	A	20010109	JP 1999-176952	19990623
JP 3550646	B2	20040804		
PRIORITY APPLN. INFO.:			JP 1999-176952	19990623
OTHER SOURCE(S):	MARPAT 134:72032			
GI				



AB The intermediates are rotaxanes of I type compds. (A, B = arom. hydrocarbyl;
 Q, R = H, C1-5 aliphatic hydrocarbyl, aromatic hydrocarbyl, hydrocarbyl
 sulfide; X = CH₂, O, S, CO, COO, OCO or CONH; n = 2-18; m = 1-5) which can be
 used as intermediates or monomers for polymers. Thus, heating 3.1 g
 diphenylacetyl-glycylglycine Et ester with 3.6 g 10-(3,5-di-tert-
 butylphenoxy)decan-1-ol in the presence of Sn catalyst in PhMe gave a dumb
 bell-like compound, 1.0 g of which was reacted with 3.1 g 5-
 allyloxyisophthalic acid dichloride and 1.6 g p-xylenediamine in CHCl₃ to give
 a rotaxane at 40% yield.

IT 315199-49-6P

RL: IMF (Industrial manufacture); PREP (Preparation)

(intermediates for manufacture of polymers containing rotaxanes having a
 cyclotetraamide ring in main chains)

RN 315199-49-6 CAPLUS

CN L-Methionine, N-(diphenylacetyl)glycyl-, 10-[3,5-bis(1,1-
 dimethylethyl)phenoxy]decyl ester, rotaxane compd. with
 7,22-bis(2-propenyloxy)-3,11,18,26-tetraazapentacyclo[26.2.2.2¹³,16.15,9.1


20,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 315199-37-2

CMF C45 H64 N2 O5 S


Absolute stereochemistry.



CM 2

CRN 169203-96-7

CMF C38 H36 N4 O6



L6 ANSWER 7 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:363906 CAPLUS Full-text

DOCUMENT NUMBER: 133:79887

TITLE: Self-assembly of mechanically interlocked and threaded rings. A HREELS and XPS study of thiol-functionalized catenane and rotaxane molecules on Au(111)

AUTHOR(S): De Nadai, C.; Whelan, C. M.; Perollier, C.; Clarkson, G.; Leigh, D. A.; Caudano, R.; Rudolf, P.

CORPORATE SOURCE: Laboratoire Interdisciplinaire de Spectroscopie Electronique, Facultes Universitaires Notre-Dame de la Paix, Namur, B-5000, Belg.

SOURCE: Surface Science (2000), 454-456, 112-117

CODEN: SUSCAS; ISSN: 0039-6028

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Thiol-functionalized catenane and rotaxane thin films were investigated to understand the self-assembly of such complex mols. on Au(111). Adsorption from the liquid phase at 300 K leads to the formation of overlayers without long-range order, as evidenced by high-resolution electron energy-loss spectroscopy (HREELS). As expected for thiol adsorption, the S 2p binding

energies determined by XPS are consistent with the formation of thiolate surface intermediates. The properties of these films are explored as a function of annealing. Changes in vibrational spectra such as the emergence of a Au-O band and variations in core-level binding energies and intensities reveal mol. rearrangement due to partial desorption. In addition, based on coherent domain sizes estimated by the angular width of the elastic beam intensity, annealing promotes long-range order within the adlayers.

IT 280129-95-5

RL: PEP (Physical, engineering or chemical process); PRP (Properties); PROC (Process)

(rotaxane; self-assembly of mech. interlocked and threaded rings of thiol-functionalized catenane and rotaxane mols. on Au(111) studied by HREELS and XPS)


RN 280129-95-5 CAPLUS

CN Glycine, N-[4-[(10-mercaptodecyl)oxy]-3,5-dimethoxybenzoyl]glycyl-, 2,2-diphenylethyl ester, rotaxane compd. with 3,11,18,26-tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 280129-94-4


CMF C37 H48 N2 O7 S



CM 2

CRN 169203-75-2

CMF C32 H28 N4 O4



REFERENCE COUNT:

16

THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L6 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1997:727264 CAPLUS Full-text

DOCUMENT NUMBER: 127:293615

TITLE: Peptide-Based Molecular Shuttles

AUTHOR(S): Lane, Alexander S.; Leigh, David A.; Murphy, Aden
 CORPORATE SOURCE: Department of Chemistry, University of Manchester
 Institute of Science and Technology, Manchester, M60
 1QD, UK
 SOURCE: Journal of the American Chemical Society (1997),
 119(45), 11092-11093
 CODEN: JACSAT; ISSN: 0002-7863
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



I

AB Peptide-based mol. shuttles, composed of macrocycle I threaded on
 glycyglycine chains [Ph₂CHCO-Gly-Gly-O(CH₂)₃]₂X [X = CH₂CH₂, (CH₂)₁₀, S],
 were prepared and their solvent-dependent translational isomerism studied by
¹H NMR. Calcns. based on variable temperature ¹H NMR data give a ΔG_{thermod.}
 for shuttling in halogenated solvents (CDCl₃, CD₂Cl₂) of 11.2, 12.4, and and
 10.9 ± 0.3 kcal mol⁻¹, resp., for I, which correspond to shuttling rates of
 37,000, 5200, and 62,000 s⁻¹, resp., at 298K. Chemical derivatization of I (X
 = S) to the corresponding sulfoxide or sulfone has no effect on the rate of
 shuttling, but the shuttling can be stopped by introduction of a bulky
 tosylimino group.

IT 196867-86-4P

RL: PEP (Physical, engineering or chemical process); PRP (Properties); SPN
 (Synthetic preparation); PREP (Preparation); PROC (Process)
 (preparation and positional isomerization of peptide-based rotaxane mol.
 shuttles)


RN 196867-86-4 CAPLUS

CN Glycine, N-(diphenylacetyl)glycyl-, [[(4-methylphenyl)sulfonyl]sulfinimido
 yl]di-3,1-propanediyl ester, rotaxane compd. with 3,11,18,26-
 tetraazapentacyclo[26.2.2.2.13,16.15,9.120,24]hexatriaconta-
 5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone
 (1:1) (9CI) (CA INDEX NAME)


CM 1

CRN 196867-85-3

CMF C49 H53 N5 O10 S2



CRN 169203-75-2
CMF C32 H28 N4 O4



=> FIL REGISTRY

FILE 'REGISTRY' ENTERED AT 16:00:13 ON 28 AUG 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 American Chemical Society (ACS)

```
STRUCTURE FILE UPDATES: 27 AUG 2007 HIGHEST RN 945649-99-0
DICTIONARY FILE UPDATES: 27 AUG 2007 HIGHEST RN 945649-99-0
```

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=> S 196867-86-4/RN

L7 1 196867-86-4/RN

=> SET NOTICE 1 DISPLAY

NOTICE SET TO 1 U.S. DOLLAR FOR DISPLAY COMMAND
SET COMMAND COMPLETED

=> D L7 SQIDE 1-

YOU HAVE REQUESTED DATA FROM 1 ANSWERS - CONTINUE? Y/(N):y
THE ESTIMATED COST FOR THIS REQUEST IS 6.55 U.S. DOLLARS
DO YOU WANT TO CONTINUE WITH THIS REQUEST? (Y)/N:y

L7 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2007 ACS on STN
RN 196867-86-4 REGISTRY
CN Glycine, N-(diphenylacetyl)glycyl-, [[[4-methylphenyl)sulfonyl]sulfinimido
yl]di-3,1-propanediyl ester, rotaxane compd. with 3,11,18,26-
tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-
5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone
(1:1) (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 3,11,18,26-Tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-
5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone,
rotaxane compd. with N-(diphenylacetyl)glycylglycine [[[4-
methylphenyl)sulfonyl]sulfinimidoyl]di-3,1-propanediyl ester (1:1) (9CI)

MF C49 H53 N5 O10 S2 . C32 H28 N4 O4

SR CA

LC STN Files: CA, CAPLUS



DT.CA Caplus document type: Journal

RL.NP Roles from non-patents: PREP (Preparation); PROC (Process); PRP
(Properties)

CM 1

CRN 196867-85-3


CMF C49 H53 N5 O10 S2



CM 2

CRN 169203-75-2

CMF C32 H28 N4 O4



1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> SET NOTICE LOGIN DISPLAY

NOTICE SET TO OFF FOR DISPLAY COMMAND
SET COMMAND COMPLETED

=>

=> logoff hold


<http://www.cas.org/support/stngen/stndoc/properties.html>

=> s 169203-75-2

L1 1 169203-75-2
(169203-75-2/RN)

=> d scan

L1 1 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
IN 3,11,18,26-Tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-
5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone
(9CI)
MF C32 H28 N4 O4
CI COM



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

ALL ANSWERS HAVE BEEN SCANNED

ENTER DISPLAY FORMAT (IDE):all

L1 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2007 ACS on STN

RN 169203-75-2 REGISTRY

ED Entered STN: 20 Oct 1995

CN 3,11,18,26-Tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (9CI) (CA INDEX NAME)

OTHER NAMES:

CN 1,7,14,20-Tetraaza-2,6,15,19-tetraoxo-3,5,9,12,16,18,22,25-tetrabenzocyclohexacosane

MF C32 H28 N4 O4

CI COM

SR CA

LC STN Files: CA, CAPLUS, CASREACT, TOXCENTER

DT.CA Caplus document type: Journal; Patent


RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)

RL.NP Roles from non-patents: ANST (Analytical study); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)

RLD.NP Roles for non-specific derivatives from non-patents: RACT (Reactant or reagent)

Ring System Data

Elemental Analysis EA	Elemental Sequence ES	Size of the Rings SZ	Ring System Formula RF	Ring Identifier RID	RID Occurrence Count
=====	=====	=====	=====	=====	=====
C6-C6-C6-C6-C22N4	C6-C6-C6-C6-NC5NC6NC5NC6	6-6-6-6-26	C32N4	19506.4.5	1



Predicted Properties (PPROP)

PROPERTY (CODE)	VALUE	CONDITION	NOTE
Bioconc. Factor (BCF)	11.64	pH 1 25 deg C	(1)
Bioconc. Factor (BCF)	11.73	pH 2 25 deg C	(1)
Bioconc. Factor (BCF)	11.74	pH 3 25 deg C	(1)
Bioconc. Factor (BCF)	11.74	pH 4 25 deg C	(1)
Bioconc. Factor (BCF)	11.74	pH 5 25 deg C	(1)
Bioconc. Factor (BCF)	11.74	pH 6 25 deg C	(1)
Bioconc. Factor (BCF)	11.74	pH 7 25 deg C	(1)
Bioconc. Factor (BCF)	11.74	pH 8 25 deg C	(1)
Bioconc. Factor (BCF)	11.74	pH 9 25 deg C	(1)
Bioconc. Factor (BCF)	11.74	pH 10 25 deg C	(1)
Boiling Point (BP)	1028.5+/-65.0 deg C	760 Torr	(1)
Density (DEN)	1.199+/-0.06 g/cm**3	760 Torr	(1)
Enthalpy of Vap. (Hvap)	150.58+/-3.0 kJ/mol	760 Torr	(1)
Flash Point (FP)	316.0+/-34.4 deg C		(1)
Freely Rotatable Bonds (FRB)	0		(1)
H acceptors (HAC)	8		(1)
H donors (HD)	4		(1)
Hydrogen Donors/Acceptors Sum (HDAS)	12		(1)
Koc (KOC)	201.09	pH 1 25 deg C	(1)
Koc (KOC)	202.62	pH 2 25 deg C	(1)
Koc (KOC)	202.78	pH 3 25 deg C	(1)
Koc (KOC)	202.79	pH 4 25 deg C	(1)
Koc (KOC)	202.79	pH 5 25 deg C	(1)
Koc (KOC)	202.79	pH 6 25 deg C	(1)
Koc (KOC)	202.79	pH 7 25 deg C	(1)
Koc (KOC)	202.79	pH 8 25 deg C	(1)
Koc (KOC)	202.79	pH 9 25 deg C	(1)
Koc (KOC)	202.75	pH 10 25 deg C	(1)
LOGD (LOGD)	1.71	pH 1 25 deg C	(1)
LOGD (LOGD)	1.71	pH 2 25 deg C	(1)
LOGD (LOGD)	1.71	pH 3 25 deg C	(1)
LOGD (LOGD)	1.71	pH 4 25 deg C	(1)
LOGD (LOGD)	1.71	pH 5 25 deg C	(1)
LOGD (LOGD)	1.71	pH 6 25 deg C	(1)
LOGD (LOGD)	1.71	pH 7 25 deg C	(1)
LOGD (LOGD)	1.71	pH 8 25 deg C	(1)
LOGD (LOGD)	1.71	pH 9 25 deg C	(1)
LOGD (LOGD)	1.71	pH 10 25 deg C	(1)
LOGP (LOGP)	1.710+/-0.678	25 deg C	(1)
Mass Intrinsic Solubility	0.0038 g/L	25 deg C	(1)

(ISLB.MASS)			
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 1 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 2 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 3 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 4 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 5 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 6 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 7 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 8 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 9 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	pH 10 25 deg C	(1)
Mass Solubility (SLB.MASS)	0.0038 g/L	Unbuffered Water	(1)
		pH 7.00	
		25 deg C	
Molar Intrinsic Solubility (ISLB.MOL)	0.0000071 mol/L	25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 1 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 2 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 3 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 4 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 5 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 6 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 7 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 8 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 9 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	pH 10 25 deg C	(1)
Molar Solubility (SLB.MOL)	0.0000071 mol/L	Unbuffered Water	(1)
		pH 7.00	
		25 deg C	
Molar Volume (MVOL)	444.1+/-3.0 cm**3/mol	20 deg C	(1)
		760 Torr	
Molecular Weight (MW)	532.59		(1)
PKA (PKA)	13.10+/-0.20	Most Acidic	(1)
		25 deg C	
PKA (PKA)	-1.07+/-0.20	Most Basic	(1)
		25 deg C	
Polar Surface Area (PSA)	116.40 A**2		(1)
Vapor Pressure (VP)	0 Torr	25 deg C	(1)

(1) Calculated using Advanced Chemistry Development (ACD/Labs) Software V8.14
((C) 1994-2007 ACD/Labs)

See HELP PROPERTIES for information about property data sources in REGISTRY.

41 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

41 REFERENCES IN FILE CAPLUS (1907 TO DATE)

REFERENCE 1

AN 145:404775 CA Full-text
TI Adsorption of Fumaramide [2]Rotaxane and Its Components on a Solid
Substrate: A Coverage-Dependent Study
AU Whelan, Caroline M.; Gatti, Francesco; Leigh, David A.; Rapino, Stefania;
Zerbetto, Francesco; Rudolf, Petra
CS Laboratoire Interdisciplinaire de Spectroscopie Electronique, Facultes
Universitaires Notre-Dame de la Paix, Namur, B-5000, Belg.
SO Journal of Physical Chemistry B (2006), 110(34), 17076-17081
CODEN: JPCBFK; ISSN: 1520-6106
PB American Chemical Society

DT Journal
 LA English
 CC 66-3 (Surface Chemistry and Colloids)
 AB The coverage-dependent adsorption on Au(111) of a fumaramide [2]rotaxane and its components, a benzylic amide macrocycle and a fumaramide thread, is studied using high-resolution electron energy loss spectroscopy (HREELS). Up to monolayer coverage, the relative intensity of out-of-plane to in-plane Ph ring vibrational modes indicates that the macrocycle adopts an orientation with the Ph rings largely parallel to the surface. The formation of a chemisorption bond is evidenced by the presence of a Au-O stretching vibration. In contrast, the thread shows no evidence of chemisorption or a preferential orientation. The introduction of the thread into the macrocycle partly disrupts the film order so that the resulting chemisorbed rotaxane shows intermediate behavior with a preferential orientation up to 0.5 ML coverage. A decrease in film order and the absence of a preferred mol. orientation is observed for all three mols. at multilayer coverages. The spectral differences are addressed by mol. dynamics simulations in terms of the mobility of the phenyls of the three mols. on Au(111).
 ST adsorption fumaramide rotaxane component gold surface chemisorption
 IT Chemisorption
 Molecular orientation
 (adsorption of fumaramide rotaxane and its component on gold surface)
 IT 7440-57-5, Gold, processes
 RL: CPS (Chemical process); NUU (Other use, unclassified); PEP (Physical, engineering or chemical process); PROC (Process); USES (Uses)
 (adsorption of fumaramide rotaxane and its component on gold surface)
 IT 169203-75-2 299433-82-2 299433-83-3
 RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); PROC (Process)
 (adsorption of fumaramide rotaxane and its component on gold surface)
 RE.CNT '85 THERE ARE 85 CITED REFERENCES AVAILABLE FOR THIS RECORD
 (1) Allinger, N; J Am Chem Soc 1989, V111, P8551 CAPLUS
 (2) Altieri, A; J Am Chem Soc 2003, V125, P8644 CAPLUS
 (3) Amabilino, D; Chem Rev 1995, V95, P2725 CAPLUS
 (4) Ashton, P; Chem Eur J 2000, V6, P3558 CAPLUS
 (5) Ashton, P; Chem Eur J 2000, V6, P3558 CAPLUS
 (6) Ashton, P; J Am Chem Soc 1998, V120, P11932 CAPLUS
 (7) Badjic, J; Science 2004, V303, P1845 CAPLUS
 (8) Ballardini, R; Eur J Org Chem 2000, P591 CAPLUS
 (9) Balzani, V; Angew Chem, Int Ed 2000, V39, P3349
 (10) Balzani, V; Molecular Devices and Machines A Journey into the Nanoworld 2003
 (11) Barbara, B; Acc Chem Res 2001, V34, P409
 (12) Barbara, B; Special issue Molecular Machines 2000
 (13) Baxter, R; ChemPhysChem 2004, V5, P245 CAPLUS
 (14) Baxter, R; Langmuir 2003, V19, P7335 CAPLUS
 (15) Bermudez, V; Nature 2000, V406, P608 CAPLUS
 (16) Bissell, R; Nature 1994, V369, P133 CAPLUS
 (17) Bottari, G; Angew Chem, Int Ed 2003, V42, P5886 CAPLUS
 (18) Bottari, G; ChemPhysChem 2002, V3, P1038 CAPLUS
 (19) Cecchet, F; J Phys Chem B 2004, V108, P15192 CAPLUS
 (20) Chen, Y; Appl Phys Lett 2003, V82, P1610 CAPLUS
 (21) Chtaib, M; J Electron Spectrosc Relat Phenom 1983, V29, P293 CAPLUS
 (22) Chtaib, M; Surf Sci 1985, V162, P245 CAPLUS
 (23) Collier, C; Science 1999, V285, P391 CAPLUS
 (24) Collin, J; New J Chem 1997, P525 CAPLUS
 (25) Coronado, E; Inorg Chem 2003, V42, P6959 CAPLUS
 (26) Da Ross, T; Org Lett 2003, V5, P689
 (27) de Nadai, C; Surf Sci 2000, V454, P112
 (28) Dudek, M; Comput Chem 1995, V16, P79

- (29) Ercolessi, F; Philos Mag A 1988, V58, P213 CAPLUS
- (30) Flood, A; Aust J Chem 2004, V57, P301 CAPLUS
- (31) Flood, A; Chem Eur J 2004, V10, P6558 CAPLUS
- (32) Flores, C; Surf Sci 1990, V239, P156 CAPLUS
- (33) Fustin, C; Surf Sci 2001, V474, P37 CAPLUS
- (34) Fustin, C; Thin Solid Films 1998, V327-329, P321 CAPLUS
- (35) Gatti, F; J Am Chem Soc 2001, V123, P5983 CAPLUS
- (36) Gatti, F; Proc Natl Acad Sci U S A 2003, V100, P10 CAPLUS
- (37) Gong, C; Angew Chem, Int Ed Engl 1997, V36, P2331 CAPLUS
- (38) Hannam, J; Angew Chem, Int Ed 2004, V43, P3260 CAPLUS
- (39) Huang, T; Appl Phys Lett 2004, V85, P5391 CAPLUS
- (40) Huang, T; Nano Lett 2004, V4, P2065 CAPLUS
- (41) Ibach, H; Electron Energy Loss Spectroscopy and Surface Vibrations 1982
- (42) Jang, S; J Am Chem Soc 2005, V127, P1563 CAPLUS
- (43) Jang, Y; J Am Chem Soc 2005, V127, P4959 CAPLUS
- (44) Jeppesen, J; Eur J Org Chem 2005, P196
- (45) Jimenez, M; Angew Chem, Int Ed 2000, V39, P3284 CAPLUS
- (46) Jun, S; Tetrahedron Lett 2000, V41, P471 CAPLUS
- (47) Katz, E; J Am Chem Soc 2004, V126, P15520 CAPLUS
- (48) Kay, E; Functional Artificial Receptors 2005, P333 CAPLUS
- (49) Keaveney, C; Angew Chem, Int Ed 2004, V43, P1222 CAPLUS
- (50) Kidd, T; Angew Chem, Int Ed 2003, V42, P3379 CAPLUS
- (51) Kihara, N; Org Lett 2004, V6, P1693 CAPLUS
- (52) Kundrot, C; Comput Chem 1991, V12, P402 CAPLUS
- (53) Lane, A; J Am Chem Soc 1997, V119, P11092 CAPLUS
- (54) Laursen, B; Org Lett 2004, V6, P4167 CAPLUS
- (55) Lee, I; Langmuir 2004, V20, P5809 CAPLUS
- (56) Lee, J; Chem Commun 2001, P1042
- (57) Lee, K; Surf Sci 1999, V420, P1115 CAPLUS
- (58) Leigh, D; Chem Commun 2004, P2262 CAPLUS
- (59) Leng, Y; J Phys Chem B 2003, V107, P11940 CAPLUS
- (60) Leon, S; Chem Eur J 2002, V8, P4854 CAPLUS
- (61) Long, B; J Am Chem Soc 2003, V125, P15490 CAPLUS
- (62) Long, B; J Am Chem Soc 2003, V125, P15490 CAPLUS
- (63) Martinez-Diaz, M; Angew Chem, Int Ed Engl 1997, V36, P1904 CAPLUS
- (64) Mendoza, S; J Chem Phys 2005, V123, P244708
- (65) Montalti, M; Langmuir 2003, V19, P5172 CAPLUS
- (66) Nikitin, K; J Am Chem Soc 2005, V127, P8067 CAPLUS
- (67) Norgaard, K; J Phys Chem B 2005, V109, P1063 CAPLUS
- (68) Ponder, J; Comput Chem 1987, V8, P1016 CAPLUS
- (69) Rapino, S; Langmuir 2005, V21, P2512 CAPLUS
- (70) Rappe, A; J Phys Chem 1991, V95, P3358 CAPLUS
- (71) Sadreev, A; J Chem Phys 2004, V120, P954 CAPLUS
- (72) Sauvage, J; Molecular Catenanes, Rotaxanes and Knots 1999
- (73) Schill, G; Catenanes, Rotaxanes and Knots 1971
- (74) Steuerman, D; Angew Chem, Int Ed 2004, V43, P6486 CAPLUS
- (75) Takata, T; Polym Synth 2004, V171, P1 CAPLUS
- (76) Teobaldi, G; J Am Chem Soc 2003, V125, P7388 CAPLUS
- (77) Thiry, P; Phys Magn 1981, V4, P35 CAPLUS
- (78) Tseng, H; Angew Chem, Int Ed 2003, V42, P1491 CAPLUS
- (79) Tseng, H; Chem Eur J 2004, V10, P155 CAPLUS
- (80) Tseng, H; ChemPhysChem 2004, V5, P111 CAPLUS
- (81) Vance, A; Nano Lett 2003, V3, P81 CAPLUS
- (82) Weber, N; Inorg Chem 2003, V42, P6780 CAPLUS
- (83) Whelan, C; J Phys Chem 2002, V106, P8739 CAPLUS
- (84) Whelan, C; Surf Sci 2001, V474, P71 CAPLUS
- (85) Yoon, B; J Phys Chem B 2003, V107, P5882 CAPLUS

AN 145:15410 CA Full-text

TI Surface Enhanced Second Harmonic Generation from Macrocycle, Catenane, and Rotaxane Thin Films: Experiments and Theory

AU Arfaoui, Imad; Bermudez, Veronica; Bottari, Giovanni; De Nadai, Celine; Jalkanen, Jukka-Pekka; Kajzar, Francois; Leigh, David A.; Lubomska, Monika; Mendoza, Sandra M.; Niziol, Jacek; Rudolf, Petra; Zerbetto, Francesco

CS Materials Science Centre, University of Groningen, Groningen, NL-9747 AG, Neth.

SO Journal of Physical Chemistry B (2006), 110(15), 7648-7652
CODEN: JPCBFK; ISSN: 1520-6106

PB American Chemical Society

DT Journal

LA English

CC 66-3 (Surface Chemistry and Colloids)

AB Surface enhanced second harmonic generation (SE SHG) expts. on mol. structures, macrocycles, catenanes, and rotaxanes, deposited as monolayers and multilayers by vacuum sublimation on silver, are reported. The measurements show that the mols. form ordered thin films, where the highest degree of order is observed in the case of macrocycle monolayers and the lowest in the case of rotaxane multilayers. The second harmonic generation activity is interpreted in terms of elec. field induced second harmonic (EFISH) generation where the elec. field is created by the substrate silver atoms. The measured second order nonlinear optical susceptibility for a rotaxane thin film is compared with that obtained by considering only EFISH contribution to SHG intensity. The elec. field on the surface of a silver layer is calculated by using the Delphi4 program for structures obtained with TINKER mol. mechanics/dynamics simulations. An excellent agreement is observed between the calculated and the measured SHG susceptibilities.

ST surface enhanced second harmonic generation macrocycle catenane rotaxane film

IT Monolayers
Second-harmonic generation
(surface-enhanced second harmonic generation of macrocycle, catenane, and rotaxane film on silver)

IT 7440-22-4, Silver, uses
RL: NUU (Other use, unclassified); USES (Uses)
(surface-enhanced second harmonic generation of macrocycle, catenane, and rotaxane film on silver)

IT 169203-75-2 252266-63-0 299433-83-3
RL: PRP (Properties)
(surface-enhanced second harmonic generation of macrocycle, catenane, and rotaxane film on silver)

RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD

- (1) Allinger, N; J Am Chem Soc 1989, V111, P8551 CAPLUS
- (2) Amabilino, D; Chem Rev 1995, V95, P2725 CAPLUS
- (3) Bermudez, V; Opt Mat 2002, V21, P39
- (4) Choy, M; Phys Rev B 1976, V14, P16993
- (5) Dudek, M; J Comput Chem 1995, V16, P791 CAPLUS
- (6) Fustin, C; Surf Sci 2001, V474, P37 CAPLUS
- (7) Fustin, C; Thin Solid Films 1998, V327-329, P321 CAPLUS
- (8) Gase, T; Adv Mater 1999, V11, P1303 CAPLUS
- (9) Gase, T; Nonlinear Opt 1999, V22, P491 CAPLUS
- (10) Gatti, F; J Am Chem Soc 2001, V123, P5983 CAPLUS
- (11) Gellert, M; Annu Rev Biochem 1981, V50, P879 CAPLUS
- (12) Jalkanen, J; J Phys Chem In press
- (13) Johnston, A; Angew Chem, Int Ed Engl 1995, V34, P1209 CAPLUS
- (14) Johnston, A; Angew Chem, Int Ed Engl 1995, V34, P1212 CAPLUS
- (15) Johnston, A; J Am Chem Soc 1996, V118, P10662 CAPLUS
- (16) Kundrot, C; J Comput Chem 1991, V12, P402 CAPLUS

- (17) Lane, A; J Am Chem Soc 1997, V119, P11092 CAPLUS
- (18) Leigh, D; Nature 2003, V424, P174 CAPLUS
- (19) Mendoza, S; J Chem Phys 2005, V123, P244708
- (20) Murakami, H; J Am Chem Soc 1997, V119, P7605 CAPLUS
- (21) Ponder, J; J Comput Chem 1987, V8, P1016 CAPLUS
- (22) Rappe, A; J Phys Chem 1991, V95, P3358 CAPLUS
- (23) Raymo, F; Templated Organic Synthesis 2000
- (24) Rocchia, W; J Comput Chem 2002, V23, P128 CAPLUS
- (25) Rocchia, W; J Phys Chem B 2001, V105, P6507 CAPLUS
- (26) Sauvage, J; Molecular catenanes, rotaxanes and knots 1999
- (27) Sutton, A; Philos Mag Lett 1990, V61, P139
- (28) Wasserman, E; J Am Chem Soc 1960, V82, P4433 CAPLUS
- (29) Whelan, C; J Phys Chem B 2002, V106, P8739 CAPLUS
- (30) Whelan, C; Surf Sci 2001, V474, P71 CAPLUS

REFERENCE 3

AN 144:179928 CA Full-text

TI Surface enhanced SHG from macrocycle, catenane and rotaxane thin films: experiments and theory

AU Arfaoui, Imad; Bermudez, Veronika; De Nadai, Celine; Jalkanen, Jukka-Pekka; Kajzar, Francois; Leigh, David; Lubomska, Monika; Mendoza, Sandra M.; Niziol, Jacek; Rudolf, Petra; Zerbetto, Francesco

CS Mater. Sci. Cent., Univ. of Groningen, Groningen, NL-9747, Neth.

SO Proceedings of SPIE-The International Society for Optical Engineering (2005), 5724 (Organic Photonic Materials and Devices VII), 139-148
CODEN: PSISDG; ISSN: 0277-786X

PB SPIE-The International Society for Optical Engineering

DT Journal

LA English

CC 73-10 (Optical, Electron, and Mass Spectroscopy and Other Related Properties)

AB Surface enhanced second harmonic generation expts. on supramols.: macrocycles, catenanes and rotaxanes, monolayers and multilayers deposited by vacuum evaporation on silver layers are reported and described. The measurements show that the mol. are ordered in thin films. The highest order is observed in the case of macrocycles and the lowest in thin films of fumaramide [2] rotaxanes. Also a better ordering is observed in the case of monolayers. The observed second harmonic generation activity is interpreted in terms of elec. field induced second harmonic generation. The elec. field contributing to SHG signal is created by silver atoms on the surface of silver layers. The measured second order NLO susceptibilities for a fumaramide [2] rotaxane is compared with that obtained by considering only EFISH contribution to SHG intensities. The elec. field on the surface of silver layer is calculated using TINKER mol. mechanics/dynamics software and the Embedded Atom model. An excellent agreement is observed between the calculated and the measured SHG susceptibilities.

ST macrocycle catenane rotaxane film surface enhanced SHG expt theory

IT Electric field effects
Nonlinear optical susceptibility
Polarization
(surface enhanced second harmonic generation from macrocycles, catenanes, rotaxanes, monolayers and multilayers deposited by vacuum evaporation on silver and gold layers)

IT Films
Second-harmonic generation
(surface enhanced second harmonic generation from macrocycles, catenanes, rotaxanes, monolayers and multilayers deposited by vacuum evaporation on silver layers)

IT 7440-57-5, Gold, uses

RL: NUU (Other use, unclassified); USES (Uses)
(surface enhanced second harmonic generation from macrocycles,
catenanes, rotaxanes, monolayers and multilayers deposited by vacuum
evaporation on silver and gold layers)

IT 7440-22-4, Silver, uses

RL: NUU (Other use, unclassified); USES (Uses)
(surface enhanced second harmonic generation from macrocycles,
catenanes, rotaxanes, monolayers and multilayers deposited by vacuum
evaporation on silver layers)

IT 169203-75-2 184017-69-4 184017-75-2

RL: PRP (Properties)
(surface enhanced second harmonic generation from macrocycles,
catenanes, rotaxanes, monolayers and multilayers deposited by vacuum
evaporation on silver layers)

RE.CNT 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD

- (1) Allinger, N; J Am Chem Soc 1989, V111, P8551 CAPLUS
- (2) Bermudez, V; Opt Mat 2002, V21, P39
- (3) Choy, M; Phys Rev 1976, VB14, P16993
- (4) Dudek, M; J Comput Chem 1995, V16, P791 CAPLUS
- (5) Fustin, C; Surface Science 2001, V474, P37 CAPLUS
- (6) Fustin, C; Thin Solid Films 1998, V327-329, P321 CAPLUS
- (7) Gase, T; Adv Mater 1999, V11(15), P1303 CAPLUS
- (8) Gase, T; Nonlinear Opt 1999, V22, P491 CAPLUS
- (9) Gatti, F; J Am Chem Soc 2001, V123, P5983 CAPLUS
- (10) Johnston, A; Angew Chem Int Ed Engl 1995, V34, P1209 CAPLUS
- (11) Johnston, A; Angew Chem Int Ed Engl 1995, V34, P1212 CAPLUS
- (12) Johnston, A; J Am Chem Soc 1996, V118, P10662 CAPLUS
- (13) Kundrot, C; J Comput Chem 1991, V12, P402 CAPLUS
- (14) Lane, A; J Am Chem Soc 1997, V119, P11092 CAPLUS
- (15) Ponder, J; J Comput Chem 1987, V8, P1016 CAPLUS
- (16) Rappe, A; J Phys Chem 1991, V95, P3358 CAPLUS
- (17) Rocchia, W; J Phys Chem B 2001, V105, P6507 CAPLUS
- (18) Whelan, M; J Phys Chem B 2002, V106, P8739

REFERENCE 4

AN 143:279363 CA Full-text

TI Antitumor agents containing an amide[2]catenane.

IN Ono, Nobufumi

PA Japan

SO Jpn. Kokai Tokkyo Koho, 7 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM C07D257-10

ICS A61K031-395; A61P035-00

CC 1-6 (Pharmacology)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 2005247790	A	20050915	JP 2004-63370	20040308
	JP 3741706	B2	20060201		
	WO 2005085215	A1	20050915	WO 2004-JP12392	20040827
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,				

AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,
 SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,
 SN, TD, TG

EP 1724265 A1 20061122 EP 2004-772348 20040827


R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
 IT, LI, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR

CN 1953966 A 20070425 CN 2004-80042350 20040827

PRAI JP 2004-63370 20040308

WO 2004-JP12392 20040827

GI



AB Antitumor agents, which are difficult to be effluxed from tumor cells, contain a self-catenane of 3,11,18,26-Tetraazapentacyclo[26.2.2.213,16.15,9.120,24]hexatriaconta-5,7,9(36),13,15,20,22,24(33),28,30,31,34-dodecaene-4,10,19,25-tetrone (I). Thus, a DMSO solution of I was electroporated to colon-26 cells to inhibit cell growth. I at 20 mg/kg p.o. showed slight toxicity on mice.

ST amide catenane antitumor electroporation

IT Antitumor agents

Electroporation

Neoplasm

(antitumor agents containing an amide[2]catenane which is slightly effluxed from tumor cells)

IT 169203-75-2

RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (self-catenane; antitumor agents containing an amide[2]catenane which is slightly effluxed from tumor cells)

REFERENCE 5

AN 142:488084 CA Full-text

TI IRAS investigation of a catenane adsorption on Au(111)

AU Fustin, C. A.; Haq, S.; Wingen, A.; Gregoire, C.; Raval, R.; Dumas, P.; Hannam, J. S.; Leigh, D. A.; Rudolf, P.

CS Laboratoire Interdisciplinaire de Spectroscopie Electronique, Facultes Universitaires Notre Dame de la Paix, Namur, B-5000, Belg.

SO Surface Science (2005), 580(1-3), 57-62

CODEN: SUSCAS; ISSN: 0039-6028

PB Elsevier B.V.

DT Journal

LA English

CC 66-3 (Surface Chemistry and Colloids)

AB The adsorption of a benzylic amide [2]catenane on a Au(1 1 1) single crystal was studied using IR reflection absorption spectroscopy. The evolution of the IR spectra during catenane deposition at different temps. of the Au substrate (300 and 90 K) was compared. Evidence of the catenane deformation upon chemisorption were found at 300 K only. At 90 K the catenane is weakly adsorbed without any deformation.

ST catenane adsorption gold surface deformation state

IT Chemisorption
Deformation (mechanical)
Surface state
Surface structure
(catenane adsorption on Au(111))

IT 169203-75-2
RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); PROC (Process)
(catenane adsorption on Au(111))

IT 7440-57-5, Gold, properties
RL: PRP (Properties)
(catenane adsorption on Au(111))

RE.CNT 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD

- (1) Amabilino, D; J Am Chem Soc 1995, V117, P1271 CAPLUS
- (2) Anon; Molecular Catenanes, Rotaxanes and Knots 1999
- (3) Asakawa, M; Adv Mater 2000, V12, P1099 CAPLUS
- (4) Avery, N; J Elec Spectrosc Rel Phenom 1983, V29, P233 CAPLUS
- (5) Ballardini, R; Acc Chem Res 2001, V34, P445 CAPLUS
- (6) Balzani, V; Angew Chem Int Ed Engl 2000, V39, P3349
- (7) Balzani, V; J Am Chem Soc 2000, V122, P3542 CAPLUS
- (8) Bellamy, L; Advances in Infrared Group Frequencies 1975, V2
- (9) Bissell, R; Nature 1994, V369, P133 CAPLUS
- (10) Cavallini, M; J Phys Chem B 2001, V105, P10826 CAPLUS
- (11) Chabal, Y; Surf Sci Rep 1988, V8, P211 CAPLUS
- (12) Collier, C; J Am Chem Soc 2001, V123, P12632 CAPLUS
- (13) Collier, C; Science 2000, V289, P1172 CAPLUS
- (14) Fanti, M; J Phys Chem A 1998, V102, P5782 CAPLUS
- (15) Fujita, H; Macromol Rapid Commun 1996, V17, P509 CAPLUS
- (16) Fujita, M; Acc Chem Res 1999, V32, P53 CAPLUS
- (17) Fujita, M; Supramol Sci 1996, V3, P37 CAPLUS
- (18) Fustin, C; Chem Phys Chem 2000, V1, P97 CAPLUS
- (19) Fustin, C; Surf Sci 2001, V474, P37 CAPLUS
- (20) Fustin, C; Surf Sci 2002, V515, P45 CAPLUS
- (21) Fustin, C; Thin Solid Films 1998, V327-329, P321 CAPLUS
- (22) Haiss, W; J Electroanal Chem 1998, V454, P107 CAPLUS
- (23) Hubin, T; Coord Chem Rev 2000, V5, P200
- (24) Johnston, A; Angew Chem Int Ed Engl 1995, V34, P1209 CAPLUS
- (25) Johnston, A; Angew Chem Int Ed Engl 1995, V34, P1212 CAPLUS
- (26) Leigh, D; Angew Chem Int Ed Engl 1996, V35, P306 CAPLUS
- (27) Leigh, D; J Am Chem Soc 1998, V120, P6458 CAPLUS
- (28) Leigh, D; Nature 2003, V424, P174 CAPLUS
- (29) Nierengarten, J; J Am Chem Soc 1994, V116, P375 CAPLUS
- (30) Pease, A; Acc Chem Res 2001, V34, P433 CAPLUS
- (31) Raehm, L; Chem Eur J 2002, V8, P2153 CAPLUS
- (32) Samori, P; Chem Phys Chem 2001, V2, P461 CAPLUS
- (33) Socrates, G; Infrared Characteristic Group Frequencies 1980
- (34) Stoddart, J; Acc Chem Res 2001, V34, P410 CAPLUS
- (35) Voegtle, F; Pure Appl Chem 1996, V68, P225 CAPLUS
- (36) Vogtle, F; Angew Chem Int Ed Engl 1993, V32, P1295

REFERENCE 6

AN 142:463228 CA Full-text

TI Optical and low frequency electric field testing of the mobility of mobile parts in catenanes and rotaxanes
 AU Nowicka, K.; Chollet, P. A.; Kajzar, F.; Bottari, G.; Gatti, F. G.; Leigh, D. A.; Miniewicz, A.
 CS Commissariat a l'Energie Atomique DRT-LIST, DECS/SE2M/LCOF, CEA/Saclay, Gif sur Yvette, 91191, Fr.
 SO Nonlinear Optics, Quantum Optics (2004), 32(1-3), 175-186
 CODEN: NOQOAP; ISSN: 1543-0537
 PB Old City Publishing, Inc.
 DT Journal
 LA English
 CC 22-3 (Physical Organic Chemistry)
 AB Optical and low frequency elec. fields are used to test the mobility of sub-parts of catenanes and rotaxanes in solution and in thin films. These mols. can be adequately tailored by the mol. engineering approach in order to enhance their response to external excitation. In particular the study of the Kerr electro-optic effect in solution permits testing of the rotational mobility of mols., or their parts, under the applied low frequency elec. field. Another possible movement can be created by illuminating the photoisomerizable rotaxane thread within its absorption band. This may lead to a "clipping" type movement made possible by the trans-cis isomerization process. The results of investigation of the influence of elec. optical and low frequency fields are presented and discussed in relation to the induced mobility of different parts of these complex mols.
 ST mobility catenane rotaxane elec field
 IT Isomerization
 (cis-trans, photochem., of fumaramide/maleimide rotaxane; optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)
 IT Isomerization
 (cis-trans, thermal, of fumaramide/maleimide rotaxane; optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)
 IT Electric field
 (frequency and voltage dependence of Kerr effect; optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)
 IT Kerr effect (electrooptical)
 (in benzylic amide catenane; optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)
 IT Molecular rotation
 (of catenane; optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)
 IT Catenanes
 RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); PROC (Process)
 (optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)
 IT Rotaxanes
 RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent)
 (optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)
 IT 509146-52-5
 RL: FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)
 (cis→trans thermal isomerization; optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)
 IT 169203-75-2
 RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP

(Physical process); PROC (Process)

(self-catenane; Kerr effect; optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)

IT 299433-83-3

RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent)

(trans→cis photoisomerization; optical and low frequency elec. field testing of the mobility of mobile parts in catenanes and rotaxanes)

RE.CNT 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD

- (1) Balzani, V; Acc Chem Res 1998, V31, P405 CAPLUS
- (2) Bermudez, V; Mol Cryst Liq Cryst 2002, V374, P343 CAPLUS
- (3) Bermudez, V; Nature 2000, V406, P608 CAPLUS
- (4) Bermudez, V; Opt Materials 2002, V21, P39
- (5) Bissell, R; Nature 1994, V369, P133 CAPLUS
- (6) Brouwer, A; Science 2001, V291, P2124 CAPLUS
- (7) Buffeteau, T; Macromolecules 2001, V34, P7514 CAPLUS
- (8) Collier, C; Science 1999, V285, P391 CAPLUS
- (9) Collier, C; Science 2000, V289, P1172 CAPLUS
- (10) Deleuze, M; J Am Chem Soc 1999, V121, P2364 CAPLUS
- (11) Gase, T; Photonics Sc News 1998, V3, P22
- (12) Heath, J; Pure Appl Chem 2000, V72, P11 CAPLUS
- (13) Joachim, C; Nature 2000, V408, P541 CAPLUS
- (14) Kippelen, B; Appl Phys Lett 1996, V68, P1748 CAPLUS
- (15) Langfong, S; Molecular Electronics 1998
- (16) Leigh, D; J Am Chem Soc 1998, V120, P6458 CAPLUS
- (17) O'Konski, C; Molecular Electronics, Theory and Methods 1981, P1 CAPLUS
- (18) Pease, A; Acc Chem Res 2001, V34, P433 CAPLUS
- (19) Raymo, F; Chem Rev 1999, V99, P1643 CAPLUS
- (20) Raymo, F; Templated Organic Synthesis 2000
- (21) Sauvage, J; Acc Chem Res 1998, V31, P611 CAPLUS

REFERENCE 7

AN 139:292240 CA Full-text

TI Clays as host matrix in the synthesis of organic macrocycles

AU Georgakilas, Vasilios; Gournis, Dimitrios; Bourlinos, Athanasios B.; Karakassides, Michael A.; Petridis, Dimitrios

CS Institute of Material Science, NCSR Demokritos Ag. Paraskevi Attikis, Athens, 15310, Greece

SO Chemistry--A European Journal (2003), 9(16), 3904-3908
CODEN: CEUJED; ISSN: 0947-6539


PB Wiley-VCH Verlag GmbH & Co. KGaA

DT Journal


LA English

CC 28-23 (Heterocyclic Compounds (More Than One Hetero Atom))

GI



I



II

AB A new approach for the synthesis of amide macrocycles, based on the use of organo-clay derivs. as controlling template, is proposed as an alternative to the rotaxane method. Dications of p-xylylenediamine inserted in the clay interlayer space act as molding pillars around which neutral diamine mols. are erected via hydrogen bonding and π - π interactions to form supramol. arrays. Condensation of diamines in the supramol. arrays with diacid chlorides yields various tetramide macrocycles, e.g., I and II, in good yields. Shape, aromaticity and dimensions of the reactants are factors affecting the condensation reaction.

ST tetramide macrocycle prepn clay matrix; diacid chloride amidation clay matrix

IT Macrocyclic compounds

RL: SPN (Synthetic preparation); PREP (Preparation)

(clays as host matrix in preparation of tetramide macrocycles from diacid chlorides and diamines)

IT Cyclization

(lactamization; clays as host matrix in preparation of tetramide macrocycles

from diacid chlorides and diamines)

IT 99-63-8, Isophthaloyl chloride 124-09-4, 1,6-Hexanediamine, reactions
539-48-0, p-Xylylenediamine 2873-74-7, Glutaryl chloride 3057-45-2,
1,4-Benzenedimethanamine dihydrochloride 3739-94-4,
2,6-Pyridinedicarbonyl chloride 3857-36-1, 2,5-Thiophenedicarbonyl
chloride

RL: RCT (Reactant); RACT (Reactant or reagent)

(clays as host matrix in preparation of tetramide macrocycles from diacid chlorides and diamines)

IT 169203-75-2P 169204-03-9P 169204-04-0P 423769-56-6P 607361-01-3P
607361-03-5P 607361-06-8P 607361-08-0P

RL: SPN (Synthetic preparation); PREP (Preparation)

(clays as host matrix in preparation of tetramide macrocycles from diacid chlorides and diamines)

IT 1318-93-0, Montmorillonite, reactions

RL: RGT (Reagent); RACT (Reactant or reagent)

(sodium-rich; clays as host matrix in preparation of tetramide macrocycles from diacid chlorides and diamines)

RE.CNT 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD

(1) Amabilino, D; Chem Rev 1995, V95, P2725 CAPLUS

(2) Ballantine, J; NATO-ASI Series C 1986, V165, P197 CAPLUS

(3) Balogh, M; Organic Chemistry Using Clays 1993

(4) Bourlinos, A; J Incl Phenom Macro Chem 2001, V40, P147 CAPLUS

(5) Cloos, P; Clays Clay Miner 1972, V20, P259 CAPLUS

- (6) Cornelis, A; NATO-ASI Series C 1986, V165, P213 CAPLUS
- (7) Farmer, V; J Phys Chem 1965, V69, P683 CAPLUS
- (8) Georgakilas, V; Angew Chem 2001, V113, P4416
- (9) Georgakilas, V; Angew Chem Int Ed 2001, V40, P4286 CAPLUS
- (10) Gil, A; Catal Rev Sci Eng 2000, V42, P145 CAPLUS
- (11) Johnston, A; Angew Chem 1995, V107, P1324
- (12) Johnston, A; Angew Chem 1995, V102, P1327
- (13) Johnston, A; Angew Chem Int Ed 1995, V34, P1209 CAPLUS
- (14) Johnston, A; Angew Chem Int Ed 1995, V34, P1212 CAPLUS
- (15) Johnston, A; J Am Chem Soc 1996, V118, P10662 CAPLUS
- (16) Klopprogge, J; J Porous Mater 1998, V5, P5 CAPLUS
- (17) Konta, J; Appl Clay Sci 1995, V10, P275 CAPLUS
- (18) Lagaly, G; Solid State Ionics 1986, V22, P43 CAPLUS
- (19) Leigh, D; Angew Chem 1997, V109, P752
- (20) Leigh, D; Angew Chem Int Ed 1997, V36, P728 CAPLUS
- (21) Lin, C; ACS Symp Ser 1992, V499, P145 CAPLUS
- (22) Lin, C; Mineralogical Society Monograph 1987, 6
- (23) Ma, Y; Microp Mesop Mater 2000, V37, P243 CAPLUS
- (24) Nikalje, M; Org Prep Proced Int 2000, V32, P1 CAPLUS
- (25) Ohtsuka, K; Chem Mater 1997, V9, P2039 CAPLUS
- (26) Pinnavaia, T; Science 1983, V220, P365 CAPLUS
- (27) Rhodes, C; Clay Miner 1994, V29, P799 CAPLUS
- (28) Seel, C; Chem Eur J 2000, V6, P21 CAPLUS
- (29) Theng, B; The Chemistry of Clay-Organic Reactions 1974

REFERENCE 8

AN 137:359718 CA Full-text

TI Effect of potassium intercalation on the electronic and vibrational properties of benzylic amide [2]catenane films

AU Fustin, C. A.; Gouttebaron, R.; Caudano, R.; Rudolf, P.; Leigh, D. A.; Fanti, M.; Krug, A.; Zerbetto, F.

CS Laboratoire Interdisciplinaire de Spectroscopie Electronique, Facultes Universitaires Notre Dame de la Paix, Namur, B-5000, Belg.

SO Surface Science (2002), 515(1), 45-52
CODEN: SUSCAS; ISSN: 0039-6028

PB Elsevier Science B.V.

DT Journal

LA English

CC 73-6 (Optical, Electron, and Mass Spectroscopy and Other Related Properties)

AB The appearance of gap states in benzylic amide catenane thin films following K intercalation was studied by EELS and quantum chemical calcns. Both theory and experience find an excitation energy for transitions into these new states of .apprx.2 eV. The characteristics of these states are discussed.

ST intercalation vibration electronic transition potassium benzylic amide catenane

IT CNDO/S
Electronic structure
Intercalation
Molecular vibration
(effect of potassium intercalation on electronic and vibrational properties of benzylic amide [2]catenane films)

IT Electron energy loss spectroscopy
(high-resolution; effect of potassium intercalation on electronic and vibrational properties of benzylic amide [2]catenane films)

IT 7440-09-7, Potassium, properties
RL: MOA (Modifier or additive use); PRP (Properties); USES (Uses)
(effect of potassium intercalation on electronic and vibrational properties of benzylic amide [2]catenane films)

IT 169203-75-2

RL: PRP (Properties)

(effect of potassium intercalation on electronic and vibrational properties of benzylic amide [2]catenane films)

RE.CNT 26. THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD

- (1) Allinger, N; J Am Chem Soc 1989, V111, P8551 CAPLUS
- (2) Barker, A; Phys Rev B 1973, V8, P5418 CAPLUS
- (3) Bellamy, L; Advances in Infrared Group Frequencies 1968
- (4) Botelho do Rego, A; J Electron Spectrosc Relat Phenom 1997, V85, P81 CAPLUS
- (5) Brouwer, A; Chem Phys 1998, V238, P421 CAPLUS
- (6) Caciuffo, R; J Chem Phys 1998, V109, P11094 CAPLUS
- (7) Ceroni, P; J Phys Chem B 1999, V103, P10171 CAPLUS
- (8) Del Bene, J; J Chem Phys 1968, V48, P1807 CAPLUS
- (9) Deleuze, M; J Am Chem Soc 1999, V121, P2364 CAPLUS
- (10) Flores, C; Surf Sci 1990, V239, P156 CAPLUS
- (11) Fustin, C; Surf Sci 2001, V474, P37 CAPLUS
- (12) Fustin, C; Thin Solid Films 1998, V327-329, P321 CAPLUS
- (13) Gomes, J; J Electroanal Chem 2000, V483, P180 CAPLUS
- (14) Gunion, R; Int J Mass Spectrom Ion Process 1992, V117, P601 CAPLUS
- (15) Johnston, A; Angew Chem Int Ed 1995, V34, P1209 CAPLUS
- (16) Johnston, A; Angew Chem Int Ed 1995, V34, P1212 CAPLUS
- (17) Lee, K; Surf Sci 1999, V420, P115 CAPLUS
- (18) Lehn, J; Supramolecular Chemistry Concepts and Perspectives 1995
- (19) Leigh, D; Angew Chem Int Ed 1996, V35, P306 CAPLUS
- (20) Leigh, D; J Am Chem Soc 1998, V120, P6458 CAPLUS
- (21) Leigh, D; J Am Chem Soc 1998, V120, P6458 CAPLUS
- (22) Lii, J; J Am Chem Soc 1989, V111, P8566 CAPLUS
- (23) Lii, J; J Am Chem Soc 1989, V111, P8576 CAPLUS
- (24) Nishimoto, K; Z Physik Chem 1957, V13, P140
- (25) Schueter, M; Z Phys 1971, V247, P263 CAPLUS
- (26) Wetterer, S; Phys Chem B 1998, V102, P9266 CAPLUS

REFERENCE 9

AN 137:253550 CA Full-text

TI Adsorption of a Benzylic Amide Macrocycle on a Solid Substrate: XPS and HREELS Characterization of Thin Films Grown on Au(111)

AU Whelan, Caroline M.; Cecchet, Francesca; Baxter, Richard; Zerbetto, Francesco; Clarkson, Guy J.; Leigh, David A.; Rudolf, Petra

CS Laboratoire Interdisciplinaire de Spectroscopie Electronique, Facultes Universitaires Notre-Dame de la Paix, Namur, B-5000, Belg.

SO Journal of Physical Chemistry B (2002), 106(34), 8739-8746
CODEN: JPCBFK; ISSN: 1520-6106

PB American Chemical Society

DT Journal

LA English

CC 66-3 (Surface Chemistry and Colloids)

AB Thin films of a benzylic amide macrocycle, the common component of a wide class of mech. interlocked mols., are prepared by vapor deposition on Au(111). The films are characterized by monochromated XPS and high resolution EELS (HREELS). The relative amts. of C, nitrogen, and oxygen are consistent with the formation of intact mol. species. At monolayer coverage, the relative intensity of out-of-plane to in-plane Ph ring vibrational modes indicates that the macrocycle adopts a nearly flat-lying conformation. The formation of a chemisorption bond is evidenced by the presence of a Au-O stretching vibration and a low binding energy component in the O 1s core level region assigned to interfacial bonding. A decrease in film order and the absence of a preferred mol. orientation is observed at higher coverages. Computer modeling of the adsorption of the macrocycle on the surface rationalizes the exptl. observations.

ST chemisorption film benzylic amide macrocycle gold XPS HREELS model

IT Configuration
 (adsorbate; characterization of thin films of benzylic amide macrocycle
 adsorbed on Au(111) using XPS and HREELS)

IT Catenanes
 Rotaxanes
 RL: MSC (Miscellaneous)
 (adsorption of benzylic amide macrocycle on Au(111) studied to make
 catenanes and rotaxanes technol. reality)

IT X-ray photoelectron spectra
 (characterization of thin films of benzylic amide macrocycle adsorbed
 on Au(111) using XPS and HREELS)

IT Chemisorbed substances
 (configuration; characterization of thin films of benzylic amide
 macrocycle adsorbed on Au(111) using XPS and HREELS)

IT Electron energy loss spectroscopy
 (high-resolution, spectra; characterization of thin films of benzylic
 amide macrocycle adsorbed on Au(111) using XPS and HREELS)

IT Simulation and Modeling
 (of adsorption of benzylic amide macrocycle on Au(111))

IT 7440-57-5, Gold, properties 169203-75-2
 RL: CPS (Chemical process); PEP (Physical, engineering or chemical
 process); PRP (Properties); PROC (Process)
 (adsorption of benzylic amide macrocycle on Au(111) studied using XPS
 and HREELS)

RE.CNT 52 THERE ARE 52 CITED REFERENCES AVAILABLE FOR THIS RECORD

- (1) Allinger, N; J Am Chem Soc 1989, V111, P8551 CAPLUS
- (2) Anelli, P; J Am Chem Soc 1991, V113, P5131 CAPLUS
- (3) Balzani, V; Acc Chem Res 1998, V31, P405 CAPLUS
- (4) Baxter, R; In preparation
- (5) Beamson, G; High-Resolution XPS of Organic Polymers; The Scienta ESCA300
 Database 1992
- (6) Bedard, T; J Am Chem Soc 1995, V117, P10662 CAPLUS
- (7) Bellamy, L; The Infrared Spectra of Complex Molecules 1966
- (8) Bermudez, V; Nature 2000, V406, P608 CAPLUS
- (9) Biscarini, F; Synth Met 1999, V102, P1466 CAPLUS
- (10) Bissell, R; Nature 1994, V369, P133 CAPLUS
- (11) Chtaib, M; J Electron Spectrosc Relat Phenom 1983, V29, P293 CAPLUS
- (12) Chtaib, M; Surf Sci 1985, V162, P245 CAPLUS
- (13) Clarkson, G; In preparation
- (14) Collin, J; New J Chem 1996, V21, P525
- (15) Czanderna, A; J Vac Sci Technol A 1991, V9, P2607 CAPLUS
- (16) Davis, A; Angew Chem, Int Ed Engl 1998, V37, P909 CAPLUS
- (17) Dudek, M; J Comput Chem 1995, V16, P791 CAPLUS
- (18) Ercolessi, F; Phys Mag A 1988, V58, P213 CAPLUS
- (19) Feynman, R; Miniaturization 1961
- (20) Fustin, C; Chem Phys Chem 2000, V1, P97 CAPLUS
- (21) Fustin, C; Submitted for publication
- (22) Fustin, C; Thin Solid Films 1998, V327-329, P321 CAPLUS
- (23) Gimzewski, J; Phys World 1998, V11, P29 CAPLUS
- (24) Ibach, H; Electron Energy Loss Spectroscopy and Surface Vibrations 1982
- (25) Johnston, A; Angew Chem, Int Ed Engl 1995, V34, P1209 CAPLUS
- (26) Johnston, A; J Am Chem Soc 1996, V118, P10662 CAPLUS
- (27) Kelly, T; Angew Chem, Int Ed Engl 1997, V36, P1866 CAPLUS
- (28) Kelly, T; J Am Chem Soc 1994, V116, P3657 CAPLUS
- (29) Kelly, T; Nature 1999, V401, P150 CAPLUS
- (30) Koumura, N; Nature 1999, V401, P152 CAPLUS
- (31) Krozer, A; J Vac Sci Technol A 1997, V15, P1704 CAPLUS
- (32) Kundrot, C; J Comput Chem 1991, V12, P402 CAPLUS
- (33) Lane, A; J Am Chem Soc 1997, V119, P11092 CAPLUS

- (34) Leigh, D; J Chem Phys 2001, V114, P5006 CAPLUS
- (35) Lii, J; J Am Chem Soc 1989, V111, P8566 CAPLUS
- (36) Lii, J; J Am Chem Soc 1989, V111, P8576 CAPLUS
- (37) Mislow, K; Chemtracts:Org Chem 1989, V2, P151
- (38) Moulder, J; Handbook of Photoelectron Spectroscopy 1992
- (39) Murakami, H; J Am Chem Soc 1997, V119, P7605 CAPLUS
- (40) Netzer, F; Langmuir 1991, V7, P2544 CAPLUS
- (41) Ponder, J; J Comput Chem 1986, V8, P1016
- (42) Rappe, A; J Phys Chem 1991, V95, P3358 CAPLUS
- (43) Riga, J; Mol Phys 1977, V34, P131 CAPLUS
- (44) Saliba, N; Surf Sci 1998, V410, P270 CAPLUS
- (45) Sauvage, J; Acc Chem Res 1998, V31, P611 CAPLUS
- (46) Sauvage, J; Molecular Catenanes, Rotaxanes and Knots 1999
- (47) Socrates, G; Infrared Characteristic Group Frequencies 1980
- (48) Takamatsu, S; Catal Soc Jpn 1992, V34, P126
- (49) Thiry, P; Phys Mag 1981, V4, P35 CAPLUS
- (50) Weiss, K; J Vac Sci Technol A 1998, V16, P1017 CAPLUS
- (51) Whelan, C; In preparation
- (52) Whelan, C; Surf Sci 2001, V474, P71 CAPLUS

REFERENCE 10

AN 136:311628 CA Full-text

TI Organoclay derivatives in the synthesis of macrocycles

AU Georgakilas, Vasilios; Gournis, Dimitrios; Petridis, Dimitrios

CS Institute of Materials Science, NCSR "Demokritos" Ag., Athens, Greece

SO Angewandte Chemie, International Edition (2001), 40(22), 4286-4288
CODEN: ACIEF5; ISSN: 1433-7851

PB Wiley-VCH Verlag GmbH

DT Journal

LA English

CC 45-4 (Industrial Organic Chemicals, Leather, Fats, and Waxes)
Section cross-reference(s): 57

AB The tetraamide macrocycle was successfully synthesized from isophthaloyl dichloride and p-xylylenediamine using a sodium montmorillonite smectic clay as a template. The pillared clay was prepared from p-xylylenediamine dihydrochloride, neutral p-xylylenediamine was inserted into the pillared clay, the tetraamide macrocycle was formed by cyclocondensation of intercalated neutral p-xylylenediamine with isophthaloyl dichloride, and the product was extracted from clay.

ST tetraazatetraoxotetrabenzocyclohexacosane macrocycle synthesis
intercalated pillared montmorillonite clay cyclocondensation template

IT Clay minerals
RL: NUU (Other use, unclassified); USES (Uses)
(acidic, intercalated; organoclay derivs. as templates in the synthesis of macrocycles)

IT Macrocyclic compounds
RL: SPN (Synthetic preparation); PREP (Preparation)
(organoclay derivs. as templates in the synthesis of macrocycles)

IT Cyclocondensation reaction
(template; organoclay derivs. as templates in the synthesis of macrocycles)

IT 1318-93-0, Swy 1, uses
RL: NUU (Other use, unclassified); USES (Uses)
(organoclay derivs. as templates in the synthesis of macrocycles)

IT 3057-45-2
RL: PEP (Physical, engineering or chemical process); PYP (Physical process); PROC (Process)
(organoclay derivs. as templates in the synthesis of macrocycles)

IT 99-63-8, Isophthaloyl dichloride 539-48-0, p-Xylylenediamine

RL: RCT (Reactant); RACT (Reactant or reagent)

(organoclay derivs. as templates in the synthesis of macrocycles)

IT 121-44-8, Triethylamine, reactions

RL: RGT (Reagent); RACT (Reactant or reagent)

(organoclay derivs. as templates in the synthesis of macrocycles)

IT 169203-75-2P

RL: SPN (Synthetic preparation); PREP (Preparation)

(organoclay derivs. as templates in the synthesis of macrocycles)

RE.CNT 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD

- (1) Amabilino, D; Chem Rev 1995, V95, P2725 CAPLUS
- (2) Anelli, P; J Am Chem Soc 1992, V114, P193 CAPLUS
- (3) Cloos, P; Clays Clay Miner 1972, V20, P259 CAPLUS
- (4) Farmer, V; J Phys Chem 1965, V69, P683 CAPLUS
- (5) Grim, R; Clay Mineralogy 1968
- (6) Houk, K; J Am Chem Soc 1999, V121, P1479 CAPLUS
- (7) Johnston, A; Angew Chem 1995, V107, P1324
- (8) Johnston, A; Angew Chem 1995, V107, P1327
- (9) Johnston, A; Angew Chem Int Ed Engl 1995, V34, P1209 CAPLUS
- (10) Johnston, A; Angew Chem Int Ed Engl 1995, V34, P1212 CAPLUS
- (11) Johnston, A; J Am Chem Soc 1996, V118, P10662 CAPLUS
- (12) Lagaly, G; Solid State Ionics 1986, V22, P43 CAPLUS
- (13) Leigh, D; Angew Chem 1997, V109, P752
- (14) Leigh, D; Angew Chem Int Ed Engl 1997, V36, P728 CAPLUS
- (15) Lin, C; "Supramolecular Architecture: Synthetic Control in Thin Films and Solids", ACS Symp Ser 1992, V499, P145 CAPLUS
- (16) Newman, A; Chemistry of Clays and Clays Minerals, Mineralogical Society Monograph, No 6 1987
- (17) Pedersen, C; J Am Chem Soc 1967, V89, P2495 CAPLUS
- (18) Pinnavaia, T; Science 1983, V220, P365 CAPLUS
- (19) Raymo, F; Chem Rev 1999, V99, P1643 CAPLUS
- (20) Seel, C; Chem Eur J 2000, V6, P21 CAPLUS
- (21) Thenk, B; The Chemistry of Clay-Organic Reactions 1974
- (22) Thompson, M; J Am Chem Soc 1962, V84, P1762 CAPLUS